

Figure 1: Mass-dependent fractionation in the solar wind has been known to occur for CMEs. This plot compares the results of for the January 1997 CME with results for the solar wind related to the May 1998 CME, as well as measurements of solar energetic particles of the November 1997 events. The data for the January 1997 event are from SOHO/CELIAS/MTOF, the May 1998 solar wind data from SWIMS on ACE, and the November data from SIS on ACE. The plot shows that there was little mass fractionation for the May 1998 event, and that the isotopic composition of Mg and Si was consistent with the solar (or terrestrial) value for this time period.

## Mass Fractionation in CMEs? ACE/SWIMS/SIS




Figure 2: Comparison of the solar wind data from the May 1998 CME with solar energetic particles from the November 1997 events. Both were acquired with instruments on ACE. The SEPs observed with ACE/SIS showed a power-law enrichment with mass, while the solar wind shows no dependence at all. Although the two events are unrelated, this observation confirms expectations that the SEPs are strongly fractionated in the acceleration process and not just due to some very unusual source population in the solar wind.




Figure 3: Solar wind data for the May 1998 CME period. The vertical axis is proportional to mass per charge, while mass is plotted along the horizontal axis. Different elements are clearly separated and some isotopes are visible.




Figure 4: Mass spectrum of the May 1998 CME period. The isotopes of Mg and Si are clearly visible. The raw data were filtered using a standard signal-processing procedure, so-called Rauch-Tung-Striebel filtering. This is a variant of the better known Kalman filter. A model fit is also shown and has been fitted to the data with a maximum-likelihood estimator.




Figure 5: Error analysis for the values in table 1. Using the known instrument response and its uncertainties, a series of artificial spectra can be generated and then subjected to the same analysis as the real data. The variations in these model data are then a good measure for the uncertainties of the measured values.

**Table 1.** Values for the isotopic composition of Mg and Si and our results for the bulk plasma in the second half of the May 1998 CME. See text for discussion.

source	$^{25}\text{Mg}/^{24}\text{Mg}$	$^{26}\text{Mg}/^{24}\text{Mg}$	ref.	$^{29}\text{Si}/^{28}\text{Si}$	$^{30}\text{Si}/^{28}\text{Si}$	ref.
FSW	$12.8 \pm 1.1$	$13.8 \pm 1.2$	a	—	—	
SSW	$13.2 \pm 1.3$	$15.3 \pm 1.3$	a	—	—	
SSW	$13.0 \pm 0.7$	$13.7 \pm 1.0$	b	$5.0 \pm 0.2$	$3.39 \pm 0.2$	h
CME	$11.7 \pm 2.1$	$14.6 \pm 2.4$	c	$5.6 \pm 1.6$	$3.6 \pm 1.5$	c
SOL	12.66	13.94	d	5.06	3.36	d
SEPs	—	$15.2 \pm 1.2$	e	$16.8 \pm 16.8$	$2.6 \pm 4.6$	i
SEPs	$14.8 \pm 3.6$	$14.8 \pm 3.4$	f	—	—	
SEPs*	$25.0 \pm 19.0$	$36.0 \pm 21.0$	g	—	—	

<sup>a</sup>Bochsler *et al.* [1997].

<sup>b</sup>Kucharek *et al.* [1998].

<sup>c</sup>This work.

<sup>d</sup>Anders and Grevesse [1989].

<sup>e</sup>Selesnick *et al.* [1993].

<sup>f</sup>Mewaldt *et al.* [1984].

<sup>g</sup>Mason *et al.* [1994].

<sup>h</sup>Wimmer-Schweingruber *et al.* [1998].

<sup>i</sup>Simpson *et al.* [1983].

SSW denotes slow solar wind, FSW fast solar wind, CME coronal mass ejection SOL solar system, and SEPs solar energetic particles. Mean of asymmetric uncertainties of Mewaldt *et al.* [1984] given. \*  $^3\text{He}$  rich event

## References

- Anders, E., and N. Grevesse, Abundances of the elements: Meteoritic and solar, *Geochim. Cosmochim. Acta*, *53*, 197 – 214, 1989.
- Bochsler, P., H. Balsiger, R. Bodmer, O. Kern, T. Zurbuchen, G. Gloeckler, D. C. Hamilton, M. R. Collier, and D. Hovestadt, Limits of the efficiency of isotope fractionation processes in the solar wind derived from the magnesium isotopic composition as observed with the WIND/MASS experiment, *Phys. Chem. Earth*, *22*, 401 – 404, 1997.
- Kucharek, H., et al., Magnesium isotopic composition as observed with the MTOF sensor on the CELIAS experiment on the SOHO spacecraft, *J. Geophys. Res.*, *103*, 26,805 – 26,812, 1998.
- Mason, G. M., J. E. Mazur, and D. C. Hamilton, Heavy-ion isotopic anomalies in  $^3\text{He}$ -rich solar particle events, *Astrophys. J.*, *425*, 843 – 848, 1994.
- Mewaldt, R. A., J. D. Spalding, and E. C. Stone, A high-resolution study of the isotopes of solar flare nuclei, *Astrophys. J.*, *280*, 892 – 901, 1984.
- Selesnick, R. S., A. C. Cummings, R. A. Leske, R. A. Mewaldt, and E. C. Stone, Coronal abundances of neon and magnesium isotopes from solar energetic particles, *Astrophys. J.*, *418*, L45 – L48, 1993.
- Simpson, J. A., J. P. Wefel, and R. Zamow, Elemental and isotopic composition of solar energetic particles, in *18-th International Conference on Cosmic Rays, Bangalore, India*, vol. 10, pp. 322 – 325, Tata Inst. Fund. Res., Bombay, India, 1983.
- Wimmer-Schweingruber, R. F., P. Bochsler, O. Kern, G. Gloeckler, and D. C. Hamilton, First determination of the silicon isotopic composition of the solar wind: WIND/MASS results, *J. Geophys. Res.*, *103*, 20,621 – 20,630, 1998.